

GST-HRP

Cat. QYA04234AHRP

Background

Epitope tags are useful for the labeling and detection of proteins using immunoblotting, immunoprecipitation, and immunostaining techniques. Because of their small size, they are unlikely to affect the tagged protein's biochemical properties. Glutathione S-transferase (GST) is a widely used fusion partner, since it provides both an easily detectable Tag and a simple purification process with little effect on the biological function of the protein of interest.

Source

The antibody was affinity-purified by affinity-chromatography using specific immunogen.

Product

Each vial contains 50ug mouse IgG diluted in 100ul of PBS pH7.4, containing 1mg/ml BSA, 0.01% thiomersal and 50% glycerol. The antibody concentration is 0.5mg/ml.

Specificity

The antibody detects C-terminal, internal, and N-terminal GST-tag fusion proteins.

Applications and Suggested Working Concentration

WB: 1:5000-1:10000

Storage

Storage at -20°C. Do not aliquot the antibody. Stable for one year from the date of shipment.

Research Use

For research use only, not for use in diagnostic procedures.

GST-HRP

| | |
|-----------------------------|--|
| Catalog No. | QYA04234AHRP |
| Size. | 100ul |
| Source. | Mouse |
| Immunogen. | Synthesized peptide |
| Purification. | The antibody was affinity-purified from mouse antiserum by affinity-chromatography using specific immunogen. |
| Specificity. | The antibody detects C-terminal, interal, and N-terminal GST-tag fusion protein. |
| Formulation. | PBS, pH 7.4, containing 0.01% thiomersal and 50% Glycerol. |
| Concentration. | 0.5 mg/ml |
| Storage / Stability. | -20°C/1 year |
| Reactivity. | N/A |
| Applications. | WB |
| Dilution. | WB:1:5000-1:10000 |